

## Algebraic topology: On some results of topological group space

Taghreed Hur Majeed  
Mustansiriyah University, Iraq

### ABSTRACT

Our main interest in this work is to study on topological group and topological groupoid spaces. We give new some results of certain types for topological group which are source proper group space denoted by (SPHG-Space), for topological groupoid are source proper topological groupoid denoted by (SPHG-Space) and for H-Space  $(H, \pi, D)$  are source proper groupoid space, denoted by (SPHH-Space). The important point is to get relationship of SPHG, SPHG-Space and SPHH-Space.

**Keywords:** Group space, proper group space, topological group, topological groupoid

### Corresponding Author:

Taghreed Hur Majeed  
Department of Mathematics, Education College  
Mustansiriyah University, Baghdad, Iraq  
E-mail:taghreedmajeed@uomustansiriyah.edu.iq

### 1. Introduction

In this paper, we study SPHG-Space, SPHG-Space and SPHH-Space and they properties for this purpose, we divide this work into section: In section one, we give the definition topological group, topological groupoid and study the properties of these spaces. In section two, we provided with several proposition about the relationship of SPHG-Space, SPHG-Space and SPHH-Space respectively. Finally, propositions in this project are presented and debated.

### 2. Topological group space

In this section, we give primary concepts of this research, Let  $(H, D)$  be a topological groupoid,  $M$  be a topological Space,  $\pi: M \rightarrow D$  be a continuous map and Let  $H * M$  denote the fiber product of  $\alpha$  and  $\pi$  over  $D$ . A left action of  $H$  on  $(M, \pi, D)$  is a continuous map  $\theta^*: H * M \rightarrow M$  such that (i)  $\pi(\theta^*(h, z)) = \beta(h)$  for each  $(h, z) \in H * M$ . (ii)  $\theta^*(w(\pi(z)), z) = z$  for each  $z \in M$ . (iii)  $\theta^*(h(\theta^*(h, z))) = \theta^*(\gamma(h, \hat{h}), z)$  for each  $(h, \hat{h}) \in H * H$  and  $(h, z) \in H * M$ . the bundle  $(M, \pi, D)$  together with action  $\theta^*$  is called groupoid space and is denoted by H-space. [5], [6]. Let  $(M, \pi, D)$  be a H-space then: The subset  $H(Z) = \{\theta^*(h, z) | h \in H_{\pi(z)}\}$  is the orbit of  $z$  under  $H$ . The action of  $H$  on  $(M, \pi, D)$  is free if for each  $(h, z) \in H * M$  the relation  $\theta^*(h, z) = z$  implies his unity. The action of  $H$  on  $(M, \pi, D)$  is a transitive if for each  $(z, \acute{z}) \in M \times M$ , there is  $h \in H$  such that  $\acute{z} = \theta^*(h, z)$  [3]. Let  $(M, \pi, D)$  be a H-space then: We say that  $(M, \pi, D)$  is free H-space if the action of  $H$  on  $(M, \pi, D)$  is free. We say that  $(M, \pi, D)$  is transitive H-space if the action of  $H$  on  $(M, \pi, D)$  is transitive. [2]. A topological groupoid  $(H, D)$  is called source proper groupoid (SPHG-Space) if: The source map  $\alpha: H \rightarrow D$  is a proper. The base space  $D$  is a Hausdorff. [4], [3]. A  $\Gamma$  - Space  $M$  is called source proper group space (SPHG-Space) if: The action groupoid  $(M \times \Gamma, M)$  is SPHG-Space.  $M$  is free  $\Gamma$  - Space. [7], [1]. Let  $(H, D)$  be an SPHG-Space then the  $\alpha$  - fiber space  $H_x$  is  $SPH_x H_x$  -space for every  $x \in D$ . A H-space  $(M, \pi, D)$  is called source proper topological groupoid space (SPHH-Space) if: The action groupoid  $(H * M, M)$  is SPHG-Space.  $(M, \pi, D)$  is free and transitive H-Space.



be SPH  $\Gamma$  – Space.  $\psi$  is free and transitive continuous action of  $H$  on  $(M, \pi, M/\Gamma)$  since:  $\pi(\psi([\dot{z}, z], \dot{z})) = \pi(\theta(\dot{z}, \delta^*(z, \dot{z}))) = \pi(\dot{z}) = \beta([\dot{z}, \dot{z}])$ .  $\psi(w(\pi(Z)), Z) = \psi([\dot{z}, z], z) = \theta(z, \delta^*(z, z)) = \theta(z, e) = z$ .

$$(i) \quad \psi([\dot{z}, z], \psi([\dot{z}, \dot{z}])) = \psi([\dot{z}, z], \theta(\dot{z}, \delta^*(\dot{z}, z_1))), \theta(\dot{z}, \delta^*(z, \theta(\dot{z}, \delta^*(\dot{z}, z_1)))) \\ = \theta(\dot{z}, \delta^*(z, \dot{z})) = \psi(\gamma([\dot{z}, z], [\dot{z}, \dot{z}]), z_1)$$

Since  $\pi(z) = \pi(\dot{z})$ .  $\Psi$  is a free action since if  $\psi([\dot{z}, z], z) = Z$  then  $\theta(\dot{z}, \delta^*(\dot{z}, z)) = Z$  and  $\theta(\dot{z}, \gamma) = Z$  and  $\theta(\dot{z}, \gamma) = Z \implies \theta(\dot{z}, \gamma) = \theta(\dot{z}, \gamma) \implies \dot{z} = \dot{z}$  and then  $[\dot{z}, \dot{z}]$  is unity.  $\Psi$  is transitive action since if  $(\dot{z}, z) \in M \times M$  then  $\psi([\dot{z}, z]) = \theta(\dot{z}, \delta^*(z, z)) = \theta(\dot{z}, e) = \dot{z}$ . Now to show that  $(H * M, M)$  is SPH-groupiod: The base space  $M$  is Hausdorff.  $H * M = (\alpha \times \pi)^{-1}(\Delta M/\Gamma)$ . The source map  $\alpha$  is proper since  $(M \times M/\Gamma), M/\Gamma$  is SPH- groupiod, since  $M/\Gamma$  is compact. Hence  $\Delta M/\Gamma$  is compact in  $M/\Gamma \times M/\Gamma$  (closed subspace of compact space) and the map  $\pi: M \rightarrow M/\Gamma$  is proper. Therefore  $H * M$  is compact since  $\alpha \times \pi$  is proper map and then it's source map is proper.

**Proposition (3):** Let  $M \times M/\Gamma, M/\Gamma$  be on SPH  $\Gamma$  – space then  $M$  is SPHH $_{\pi(Z)} H_{\pi(Z)}$  –Space, for all  $z \in M$ . Where  $\pi: M \times M/\Gamma \rightarrow M/\Gamma$ .

**Proof:** The map  $\eta_z: H_{\pi(Z)} \rightarrow M, \eta_z(h) = \theta^*(h, Z)$  where  $\theta^*$  is the law of action of  $H$ -space and  $\eta_z$  is closed map since for every closed subset  $A$  of  $H_{\pi(Z)}$  then  $A$  is compact ( $G_{\pi(Z)}$  is compact) and then  $\eta_z(A)$  is compact and consequently  $\eta_z(A)$  is closed ( $M$  is Hausdorff since  $(H * M, M)$  is SPHH-space). Hence  $\eta_z$  is homeomorphism, now define  $\phi_z = M \times_{\pi(Z)} H_{\pi(Z)} \xrightarrow{\eta_z^{-1} \times I} H_{\pi(Z)} \times_{\pi(Z)} H_{\pi(Z)} \xrightarrow{\gamma} H_{\pi(Z)} \xrightarrow{\eta_z} M$ , by:  $\phi_z(\dot{z}, r) = \eta_z(\gamma(h, r))$  where  $h$  is a unique element such that  $h = \eta_z^{-1}(\dot{z})$  and  $\gamma = \gamma|_{H_{\pi(Z)} \times_{\pi(Z)} H_{\pi(Z)}}$ .  $M$  is free $_{\pi(Z)} H_{\pi(Z)}$  –Space by  $\phi_z$  since:  $\phi_z(\dot{z}, w(\pi(Z))) = \phi_z(\eta_z(h), w(\pi(Z))) = \eta_z(\gamma(h, w(\pi(Z)))) = \eta_z(\gamma(h, w(\alpha(h)))) = \eta_z(h) = \dot{z}$  where  $w(\pi(Z))$  is unity in $_{\pi(Z)} H_{\pi(Z)}$ .  $\phi_z(\dot{z}, \gamma(r_1, r_2)) = \phi_z(\eta_z(h), \gamma(r_1, r_2)) = \eta_z(\gamma(h, \gamma(r_1, r_2))) = \eta_z(\gamma(h, r_1), r_2) = \phi_z(\phi_z(\eta_z(h), r_1), r_2) = \phi_z(\phi_z(\dot{z}, r_1), r_2)$ . If  $\phi_z(\dot{z}, r) = \dot{z}$  then  $\phi_z(\eta_z(h), r) = \eta_z(h)$  ( $\eta_z$  injective) and then  $\gamma$  is unity  $\phi_z$  is continuous since it is the composition of continuous map. Now to show the action groupoid  $(M \times_{\pi(Z)} H_{\pi(Z)}, M)$  is SPHG-Space:  $M$  is Housdorff space since  $(M, \pi, M/\Gamma)$  is SPHG-Space. (a)The Source map  $\alpha_z: M \times_{\pi(Z)} H_{\pi(Z)} \rightarrow M, \alpha_z(\dot{z}, r) = \dot{z}$  is a proper map by using the following commutative diagram:




Figure 2. Diagram of Proposition (3)

Where  $\alpha_z$  is a Source map of the action groupoid  $(H_{\pi(Z)} \times_{\pi(Z)} H_{\pi(Z)})$  which is proper map and then we have  $\alpha_z$  is proper map.

**Proposition (4):**

Let  $M$  be an SPHG-Space then

$(M \times M /_{\pi(z)} H_{\pi(z)}, M /_{\pi(z)} H_{\pi(z)})$  is SPH-Space for every  $z \in M$ .

**Proof:**

$M$  is  $SPH_{\pi(z)} H_{\pi(z)}$  - Space for every  $z \in M$  we have Ehressman groupoid  $(M \times M /_{\pi(z)} H_{\pi(z)}, M /_{\pi(z)} H_{\pi(z)})$  is SPHG-Space for every  $z \in M$ .

**Proposition (5):**

Let  $(H, D)$  be an SPH- groupoid and  $(M \times M /_{\pi(z)} H_{\pi(z)})$  be SPHG-Space then the map  $\pi: M \rightarrow D$  is proper map.

**Proof:**

Consider the following commutative diagram in  $M$  as in Figure 3.




Figure 3. Diagram of Proposition (4)

In which  $\eta_z$  is an homeorphism given proposition (4.1.10) and  $\beta_{\pi(z)} = w|_{\pi(z)H_{\pi(z)}}$  is proper map since  $(H,D)$  is SPHG-Space then  $\pi \circ \eta_z$  is proper map and then  $\pi$  is proper map.

**4. Conclusion**

New findings from certain topology groups in this paper were given, the proper grouping space denoted by (SPHG-Space), the proper topological groupoid source is denoted by (SPHG space), and H-space  $(H, \pi, D)$ , the correct groupoid source denoted by groupoid source, indicates the correct topological groupoids (SPHH-Space). The point is to have SPHG, SPHG-Space and SPHH-Space associations that are mathematically important.

**5. Acknowledgement**

The author (Taghreed Hur Majeed) would be grateful to thank Mustansiriyah University ([www.mustansiriyah.edu.id](http://www.mustansiriyah.edu.id)) in Baghdad, Iraq. For it collaboration and support in the present work.

**References**

- [1] R. Brown, "Topology and Groupoids" Book surge LIC, Deganwy, United Kingdom, 2020.
- [2] M. H. Gürsoy, "General Topological Groupoids" Punjab University Journal of Mathematics, vol.53, no.2, pp. 35-49, 2021.
- [3] M. H. Gürsoy and I. Icen, "The Homomorphisms of Topological Groupoids" NOVI SAD J. MATH., vol.44, no.1, pp. 129-141, 2014.
- [4] T. H. Majeed and S. A. Mohammed, "Category of Source Proper Group Space" Zanco Journal of pure and applied Science, vol.31, no.2, pp. 6-13, 2019.

- [5] T. H. Majeed , "On Some results of Topological Groupoid" Journal of physics IOP conference Series, vol. 1003, No. 1, p. 012067, 2018.
- [6] V. Marin and J. Avila and H. Pinedo, "Isomorphism Theorems for Groupoids and Some Application" Hindawi, International Journal of Mathematics and Mathematical Sciences, vol. 2020, 2020.
- [7] M. Q. Mann'a, " Groupoid and Topological Qoutient Group" Global Journal of pure and Applied Mathematics, vol.13, no.7, pp.3173-3191, 2017.